

Chiral Symmetry in algebraic and analytic approaches

V. Vereshagin

St.Petersburg State University, St.Petersburg, Russia
and

Institute for Theor. Physics III, Univ. of Erlangen-Nuernberg, Erlangen, Germany
M. Dillig

Institute for Theor. Physics III, Univ. of Erlangen-Nuernberg, Erlangen, Germany
A. Vereshagin

St.Petersburg State University, St.Petersburg, Russia

Abstract

We compare among themselves two different methods for the derivation of results following from the requirement of polynomial boundedness of tree-level chiral amplitudes. It is shown that the results of the algebraic approach are valid also in the framework of the analytical one. This means that the system of Sum Rules and Bootstrap equations previously obtained with the help of the latter approach can be analyzed in terms of reducible representations of the unbroken Chiral group with the known structure of the mass matrix.

Novadays it is commonly realised that the Spontaneously Broken Chiral Symmetry (SBChS) plays an essential role in the hadronization mechanism provided by the large-distance forces in OCD. The smallness of the pion mass (compared to the typical hadronic scale of 1 GeV) made it possible to develop the Chiral Perturbation Theory (ChPT) [1, 2] (for recent reviews see also [3, 4, 5]) allowing one to calculate in a systematic manner the corrections to the zero-order ("bare") amplitude given by a sum of corresponding tree graphs. Thus, if the bare amplitudes are known, we can analyze the low energy processes with the arbitrary high degree of accuracy.

Unfortunately, the bare (tree-level) amplitudes themselves contain a lot of free parameters since they depend of infinite number of basically unknown coupling constants describing the off-shell hadron interactions and point-like vertices. Just because of this reason the current situation looks unsatisfactory: we have an excellent tool – ChPT – for the computation of corrections to the quantities which we know not so much about.

A remarkable step in the understanding of the bare couplings was done by S.Weinberg [6], first suggested to use the asymptotic boundedness requirements for the chiral amplitudes describing the forward scattering of massless pions on arbitrary targets (the list of the corresponding refs. can be found, e.g., in [7]; the paper [8] should be added to this list). It was shown that the SBChS – despite of its dynamical origin – manifests itself in a customary algebraic manner. This means that it is possible to classify the states with a given helicity according to linear reducible representations of unbroken Chiral group, the mass matrix being constructed as a sum of chiral scalar and 4-th component of chiral vector.

A variety of interesting results derived by many authors in the framework of Weinberg's scheme clearly demonstrates both the importance and the fruitfulness of the asymptotic requirements.

There are known two technically different approaches to the derivation of results from those requirements.

Weinberg's original approach [6] (later on called algebraic) consisting of the group theoretical analysis of mass (squared) and coupling matrices, shows both attractive features and serious limits. One of the most attractive features of this method is that it allows one to work from the very beginning with purely algebraic structures. However, it is applicable only for the case of forward scattering of massless pions; its feasibility is limited on a small number of partial waves taken into account. Moreover (as stressed by Weinberg himself [6]), from the mathematical point of view this method is only valid if the number of resonances is finite.

A different approach (first suggested in [9]) has been recently formulated in its final form in [7]. Later on we call this approach analytic. It takes advantage of the analytic structure of a tree-level amplitude, this structure being defined by the requirements of meromorphy and polynomial boundedness. In this method there is no necessity to require that pion is massless. Also, it is not restricted to forward processes only: the nonforward ones can be analyzed on the same ground. The method does not contain

any limitations on the number of resonances or partial waves included. Moreover, it follows from the results of ref. [7] that, to avoid self-contradictions in the case if a partial wave with $l \geq 2$ (or, the same, the resonance with a spin $J \geq 2$) is taken into account, one has to include also the full tower of higher spin resonances with couplings and masses strongly restricted by a certain infinite set of self-consistency (bootstrap) equations. So, compared to the algebraic approach, the analytic one looks more general, since it contains the algebraic method as a particular case. Moreover, it contains also the built-in mechanism guaranteeing the absence of mutually contradicting results in those cases when the algebraic method is applied for the simultaneous analysis of direct- and cross-channel processes.

However, as a serious drawback, the problem of algebraization of the system of bootstrap equations – which is clearly absent in the algebraic approach – looks rather complicated in the analytical framework. Thus, it is interesting to ask the question if it is possible to get Weinberg's results from the analysis of the corresponding part of a system of bootstrap equations derived with a help of the analytic method.

Below we give the positive answer to this question along with the general outline of the corresponding proof. A more detailed consideration will be published elsewhere.

Let us first recall some key points of the analytic approach (for more details see [7]). This approach is based on the following postulates:

1. The tree-level amplitude of a given binary process is a meromorphic function in a space of 3 dependent complex Mandelstam variables s, t, u ($s + t + u = \sum_{i=1}^4 m_i^2$).
2. In accordance with the crossing symmetry requirement, this amplitude describes all three channels of a process under consideration.
3. The principal part of the amplitude is completely determined by the formally written tree-level expression following from the "naive" (unitary) Feynman rules (no ghosts, no tachions!).
4. As a function of every one complex variable x (considered as the CMS energy squared in a given channel) this amplitude is polynomially bounded at zero value of the corresponding momentum transfer, the degree of a polynomial being dictated by the value of relevant Regge intercept (or, equivalently, by experiment).

The main technical tool used in the analytic approach is provided by the Cauchy method. This method allows one to construct the convergent power fraction expansion for the meromorphic function $f(z)$ of one complex variable z with a given principal part, this expansion being coordinated with the asymptotic condition of the form

$$\max_{z \in C_n} \left| \frac{f(z)}{z^{N+1}} \right| \xrightarrow{n \rightarrow \infty} 0, \quad (1)$$

where C_n denotes a smooth contour (for definiteness, a circle) which contains inside of it the first n poles ($|p_{i+1}| > |p_i|$, p_i is the i -th pole position, r_i – the corresponding residue; p_i being enumerated in the order of increasing modulo) and does not contain any other poles. The minimal integer number N in (1) shows the degree of the bounding polynomial.

With the condition (1) the Cauchy method gives:

$$f(z) = \sum_{p=0}^N f^{(p)}(0) \frac{z^p}{p!} + \sum_{i=1}^{\infty} \left(\frac{r_i}{z - p_i} - \Pi_i(z) \right). \quad (2)$$

Here $\Pi_i(z)$ stands for the so-called correcting polynomial: it is nothing but the sum of the first N terms of the power series expansion of $r_i/(z - p_i)$ around the point $z = 0$. The necessity in correcting polynomials is caused by the convergency condition. In fact, the eq. (2) provides a special case of the general theorem by Mittag-Leffler.

In a framework of the analytic method one works with the amplitude written in the form (2). In the case of forward process all the parameters appearing in (2) (namely, $f^{(0)}(0), r_i, p_i$) are just constants; otherwise, they should be considered as the functions of the momentum transfer. The form (2) is convenient because it does not contain any unwanted terms breaking the allowed asymptotic regime, thus from the very beginning there is no need to control the correctness of the asymptotics.

This is in strong contrast to the algebraic method, where results are derived from the condition of cancellation of just those "unwanted" terms which increase too rapidly with energy but appear necessarily in the scattering amplitude if the Lagrangian is written in terms of covariant derivatives (see [6]). In

this approach one calculates the lowest Laurent series expansion coefficient which breaks the allowed asymptotic regime and then requires of it to be zero.

At the first glance two approaches described above look quite different. Indeed, since the "naive" Laurent series expansion is only applied for the study of asymptotics if the number of poles is finite, it looks impossible to use it for the analysis of cross-conjugated processes where the infinite set of resonances is required to provide the correct asymptotic behavior in both channels simultaneously (see [7]). This, in turn, could mean that the results of early papers [10], based on the calculation of the Laurent coefficients, have no ground. At the same time, it is possible to show that the Sum Rules given in [10] can be also derived with a help of the form (2). So, it looks that the conclusions obtained in a framework of the algebraic method remain also valid with respect to the results of the analytic approach.

To check (and prove) this idea [11] let us consider the case $N = 0$ in (1) and (2) (just in order to simplify the corresponding formulas). In this case from the integral form of the asymptotic condition (1)

$$\oint_{C_n} \left| \frac{f(z)}{z^2} dz \right| \xrightarrow{n \rightarrow \infty} 0, \quad (3)$$

it follows that

$$\sum_{i=n+1}^{\infty} \frac{r_i}{p_i^2} \xrightarrow{n \rightarrow \infty} 0. \quad (4)$$

Next, let us consider the true Laurent series expansion

$$f(z) = \sum_{p=-\infty}^{\infty} C_p^{(n)} z^p, \quad (z \in R_n), \quad (5)$$

where R_n stands for the ring $|z| \in (|p_n|, |p_{n+1}|)$ and calculate the lowest "unwanted" coefficient $C_{+1}^{(n)}$. It is easy to get

$$C_{+1}^{(n)} = - \sum_{i=n+1}^{\infty} \frac{r_i}{p_i^2}. \quad (6)$$

To study the behavior of the amplitude $f(z)$ at large values of $|z|$ one can use the expansion (5) and take a limit $n \rightarrow \infty$. In this case – as it follows immediately from the comparison of (6) with (4) – one obtains the result

$$C_{+1}^{(n)} \xrightarrow{n \rightarrow \infty} 0, \quad (7)$$

showing that the lowest "unwanted" Laurent coefficient vanishes (along with all higher ones $C_i^{(n)}, i = +2, +3, \dots$). This result justifies Weinberg's formal method for the case of infinite number of resonances. It explains also the complete coincidence of the systems of Sum Rules derived in [10] (from the formal manipulation with Laurent series expansions) with those following directly from the convergent partial fraction expansion (2).

Summarizing, we conclude with the statement that the main result following from the algebraic method is also valid in the framework of the analytic one: in the chiral limit $m_\pi = 0$ the direct channel resonances with a given helicity fall into reducible representations of unbroken chiral group $SU_2 \times SU_2$, the mass matrix being constructed as a sum of a chiral scalar and the 4-th component of chiral vector.

This work was supported in part by DAAD (visiting Grant for V.V.), RFBR (Grant 96-02- 18017) and GRACENAS (Grant 95-06.3-13).

References

- [1] S.Weinberg. Physica, 96A (1979) 327.

- [2] J.Gasser, H.Leutwyler. *Ann. of Physics*, 158 (1984) 142.
- [3] Ulf-G. Meissner. *Rep. Progr. Phys.*, 56 (1993) 903.
- [4] A.Pich. *Rep. Progr. Phys.*, 58 (1995) 563.
- [5] J.Bijnens. *Phys. Reports*. 265 (1996) 371.
- [6] S.Weinberg. *Phys. Rev.*, 177 (1969) 2604; *Phys. Rev. Letters*, 65 (1990) 1177.
- [7] V.Vereshagin. Preprint SPbU-IP-96-20, 1996; hep-ph/9606464.
- [8] S.R.Beane. Preprint DUKE-TH-95-98, 1995; hep-ph/9512228.
- [9] V.Vereshagin. *Nucl. Phys.*, B55 (1973) 621.
- [10] A.Bolokhov, A.Manashov, M.Polyakov, V.Vereshagin. *Phys. Letters*, B303 (1993) 220; *Phys. Rev.*, D48 (1993) 3090.
- [11] In this point we follow the argumentation suggested by A.Vereshagin in his Bachelor Thesis (unpublished).